
A CHARACTERIZATION BASED ON THE ABSOLUTE DIFFERENCE OF TWO I. I. D. RANDOM VARIABLES

Prem S. Puri, et al

Purdue University Lafayette, Indiana

January 1970



This document has been approved for public release and sale.


A Characterization Based on the Absolute Difference of Two I.I.D. Random Variables

Ъу

Prem S. Puri\* and Herman Rubin\*\*

## **PURDUE UNIVERSITY**






## DEPARTMENT OF STATISTICS

### DIVISION OF MATHEMATICAL SCIENCES

CALARING HOUSE



# A Characterization Based on the Absolute Difference of Two I.I.D. Random Variables

bу

Prem S. Puri\* and Herman Rubin\*\*

Department of Statistics

Division of Mathematical Sciences

Mimeograph Series No. 218

January 1970

Distribution of this document is unlimited.

<sup>\*</sup> This investigation was supported in part by research grant GM-10525 from NIH, Public Health Service, at University of California, Berkeley.

<sup>\*\*</sup> This research was partly supported by the Office of Naval Research Contract N00014-67-A-0226-0008. Reproduction in whole or in part is permitted for any purpose of the U. S. Government.

A Characterization Based on the Absolute Difference

of Two I.I.D. Random Variables

bv

Prem S. Puri\* and Herman Rubin\*\*

Purdue University, Lafayette

- 1. INTRODUCTION. Let X<sub>1</sub> and X<sub>2</sub> be two independent and identically distributed (i.i.d.) random variables whose common distribution is same as that of a non-negative random variable X. The problem considered here is to characterize all possible distributions of X which satisfy the following property H:
- (1) H: The distributions of  $|X_1 X_2|$  and X are identical.

For instance, it is easy to verify that the discrete distribution with  $P(X=0)=P(X=a)=\frac{1}{2}$  for some positive constant a, and the exponential distribution with probability density function (p.d.f.) f where f(x)=9 exp  $(-\theta x)$ , for  $x\geq 0$ , and f(x)=0 elsewhere, with  $\theta>0$ , both satisfy the property H. The reader may find a different characterization based on  $|X_1-X_2|$  in Puri [2]. Let F denote the distribution function (D.F.) of X. It can be easily shown that if X satisfies H, the distribution of X can either be only discrete or absolutely continuous or singular and no mixture is possible. Thus one needs to consider these three possibilities separately. For the case when X is discrete let A

Distribution of this document is unlimited.

<sup>\*</sup> This investigation was supported in part by research grant GM-10525 from NIH, Public Health Service, at University of California, Berkeley.

<sup>\*\*</sup> This research was partly supported by the Office of Naval Research Contract NOCO14-67-A-0226-0008. Rep eduction in whole or in part is permitted for any purpose of the U.S. Government.

denote the set of possible discrete nonnegative values that X takes. More specifically, let

$$p_y = P(X=y), y \in A; \text{ with } \sum_{y \in A} p_y = 1.$$

It is clear that if there exists a  $y \ge 0$  with  $p_y > 0$ , then in particular A contains zero with  $p_0 > 0$ . Furthermore, from the property H, the following relations follow easily.

$$p_0 = \sum_{x \geq 0} p_x^2$$

(3) 
$$p_y = 2 \sum_{x \ge 0} p_x p_{x+y}; y > 0.$$

Similar relations are satisfied by the p.d.f. f if X satisfying H is absolutely continuous.

In section 2, we show that under H, X has a moment generating function (m.g.f.) and hence all its moments are finite. Also in theorem 1, we consider the case where X is bounded. Section 3 deals with the discrete case, and theorem 2 characterizes lattice distributions satisfying H. In section 4, we consider the absolutely continuous case. Here we study a more general question; namely if  $X_1$  and  $X_2$  are two nonnegative independent but not necessarily identically distributed random variables (r.v.) and moreover if the distributions of  $X_1$  and  $|X_1-X_2|$  are identical, then given the distribution of  $X_2$ , what can be said about the distribution of  $X_1$ ? The paper ends with a discussion in section 5, where we have few words to say about the singular case.

2. PRELIMINARY RESULTS. In the following lemma it is shown that for an X satisfying H, its m.g.f. and hence all its moments exist.

Lemma 1. The m.g.f. of a nonnegative r.v. X satisfying H exists.

<u>Proof.</u> If X satisfying H is degenerate, it is clear that P(X=0) = 1, and the lemma holds trivially. Let X be nondegenerate. Then there is a number u > 0 such that  $P(X \le u) > \frac{1}{2}$ . Using this and the property H, it follows that for every  $v \ge 0$ .

(4) 
$$P(X > v) = P(|X_1 - X_2| > v) \ge 2 P(X > u+v) P(X \le u),$$

so that

$$P(X > u+v) \le P(X > v) /2 P(X \le u),$$

for all  $v \ge 0$ . A repeated application of this leads to

$$P(X > nu) \le \left[\frac{1}{2 P(X < u)}\right]^{n-1}$$
; for  $n = 1, 2, ...$ 

From this one can easily show the existence of an  $\alpha > 0$  such that  $\mathbb{E}(\exp(\xi X))$  exists for all  $|\xi| < \alpha$ .

The following theorem provides the answer to our problem when X is bounded.

Theorem 1. Let X be nonnegative and nondegenerate. Then the following three statement. are equivalent.

- (i) X is bounded and satisfies H.
- (ii) X satisfies H and P(X=0) =  $\frac{1}{2}$ .
- (iii)  $P(X=0) = P(X=a) = \frac{1}{2}$ , for some a > 0.

<u>Proof.</u> Clearly (iii)  $\Rightarrow$  (i) and (ii). All we need to prove is that (i)  $\Rightarrow$  (iii) and (ii)  $\Rightarrow$  (iii). Let (i) hold. Since X is bounded and nondegenerate, there exist a least upper bound B > 0, such that P(X > B) = 0 and for every  $0 < c \le B$ , P(X > B - c) > 0. On the other hand since X satisfies H, we have for every such c

(5) 
$$0 < P(X > B-c) = P(|X_1-X_2| > B-c) \le 2 P(X < c) P(X > B-c),$$

which implies that for every  $0 \le \epsilon \le B$ ,  $P(X \le \epsilon) \ge \frac{1}{5}$  and in particular letting  $\epsilon$  tend to zero we have  $P(X=0) \ge \frac{1}{5}$ . This implies that X must be a discrete random

variable (r.v). Using the notation introduced in section 1 for such a case, we have

$$\frac{1}{2} \le p_0 = 7$$
  $p_x^2 < 1$ . This yields

$$p_0(1-p_0) = \sum_{x>0} p_x^2 \le {\max_{x>0} p_x} \sum_{x>0} p_x = {\max_{x>0} p_x} (1-p_0),$$

so that  $p_0 \le \max_{x>0} p_x$ . Hence we have

$$\frac{1}{2} \le p_0 \le \max_{x>0} p_x \le 1 - p_0 \le \frac{1}{2},$$

so that  $p_0 = \max_{x \ge 0} p_x = \frac{1}{2}$ , which implies that  $P(X=0) = P(X=B) = \frac{1}{2}$ . This proves that (i)  $\Rightarrow$  (iii) with B = a. Now let (ii) hold. Since  $p_0 = \frac{1}{2}$ , X must be a discrete r.v if it has to satisfy H. For this we have already seen that  $p_0(1-p_0) = \sum_{x \ge 0} p_x^2$ . This means that we have

$$\sum_{x>0} p_x^2 = \frac{1}{4} \text{ and } \sum_{x>0} p_x = \frac{1}{2}.$$

But this holds if and only if  $p_x = \frac{1}{3}$  for some x = a > 0, so that (iii) holds.

Q.E.1

Before closing this section we wish to remark that for the nondegenerate discrete case, for X satisfying H we must have  $0 < p_0 \le \frac{1}{2}$ . That  $p_0 > 0$  follows from (2) and the fact that  $\sum_{x \ge 0} p_x = 1$ . That  $p_0 \le \frac{1}{2}$  follows from the fact that

for every y > 0 with  $p_y$  > 0,  $p_y \ge 2 p_0 p_y$  under H.

- (6) C: There exists an interval  $(\delta_1, \delta_2)$  with  $0 \le \delta_1 \le \delta_2 \le \infty$  such that  $P(\delta_1 \le X \le \delta_2) = 0$ .

Using the notation of section 1, we first prove three lemmas needed to prove the main result of Theorem 2.

Lemma 2. Let X be discrete, nondegenerate and satisfy H and the condition C.

Then

(i) 
$$\tau = \inf \{x: x > 0, p_x > 0\} > 0 \text{ and } p_{\tau} > 0, \text{ and }$$

(ii) the set of possible values of X is given by  $k\tau$ ; k = 0,1,2,...

<u>Proof.</u> (i) Since X is nondegenerate, it is clear that the set  $\{x: x > 0, p_x > 0\}$  is nonemty. Again if  $p_0 = \frac{1}{2}$ , (i) and (ii) are satisfied in view of Theorem 1, so that let  $0 < p_0 < \frac{1}{2}$ . By Theorem 1, this means that X is not bounded. Let S be the set of possible values of X. In view of the property H, it is easy to show that S forms a positive linear space in integers. By this we mean that if  $x_i \in S$ ,  $i = 1, 2, \ldots$ , then  $\left|\sum_i x_i\right| \in S$  for all integer values (positive or negative) of  $n_i$ 's. Now for an unbounded set with this property, it is not difficult to show that either this set is dense everywhere over  $[0, \infty)$  or is a lattice. On the other hand in view of condition C, it cannot be dense everywhere. Hence the lemma follows.

In view of lemma 2, let  $p_k$  denote the probability  $p(X=k\tau)$ , for  $k=0,1,2,\ldots$ , so that  $\sum_{k=0}^{\infty}p_k=1$ . The analogues of (2) and (3) are given by

$$p_{0} = \sum_{i=0}^{\infty} p_{i}^{2}$$

(16) 
$$p_{k} = 2 \sum_{i=0}^{\infty} r_{i} r_{i+k}; k = 1,2,...$$

We shall now restrict to the case with  $C \leq p_C \leq \frac{1}{2}$ . Thus the set of possible values of X, in view of Theorem 1, must be infinitely denumerable. Furthermore, from (12) of the following lemma it follows that under condition C,  $p_k \geq C$ , for  $k=1,2,\ldots$ 

Lemma 3. Let H and C hold and also let  $0 < p_0 < \frac{1}{2}$ . Then for k = 1, 2, ...,

(11) 
$$p_{k} \ge \left(\frac{2 p_{1}}{1-2 p_{0}}\right) \cdot p_{k+1}$$

and

(12) 
$$p_{k+1} \ge \frac{2 p_1 (1-2p_0)}{[(1-2p_0)^2 + 4 p_1^2]} \cdot p_k$$

<u>Proof.</u> (11) follows easily from (10) by noticing that for  $k \ge 1$ ,

(13) 
$$p_k(1-2p_0) - 2p_1p_{k+1} = 2 \sum_{i=2}^{\infty} p_i p_{i+k},$$

and that the right side of (13) is nonnegative. To prove (12), we first notice from (10) that for  $k \ge 1$ ,

(14) 
$$p_{k+1} = 2 \sum_{i=0}^{\infty} p_i p_{i+k+1},$$

or equivalently

(15) 
$$p_{k+1}(1-2p_0) = 2 \sum_{i=1}^{\infty} p_i p_{i+k+1}.$$

Then using (11) for each p on the right side of (15) we have

$$(16) p_{k+1}(1-2p_0) \ge \frac{\mu_{p_1}}{1-2p_0} \sum_{i=1}^{\infty} p_{i+1} p_{i+k+1} = \frac{2p_1}{1-2p_0} (2 \sum_{i=2}^{\infty} p_{i} p_{i+k})$$

$$= \frac{2p_1}{1-2p_0} (p_k - 2p_0 p_k - 2p_1 p_{k+1}) .$$

Here at the end of (16) we have again used (10). Finally (12) follows immediately from (16) after a little simplification. For each sequence  $\{p_k\}$  satisfying H (or equivalently (9) and (10)) and with  $0 < p_0 < \frac{1}{2}$ , define

 $\beta = \sup \{b: b > 0 \text{ satisfying } p_k \ge b p_{k+1} \text{ for all } k \ge 1\}$  and

 $\gamma = \sup \{c: 0 \le c \le 1, \text{ satisfying } p_{k+1} \ge c p_k \text{ for all } k \ge 1\},$  so that

(17) 
$$p_k \ge 3 p_{k+1}$$
;  $k = 1,2,...$ 

and

(18) 
$$p_{k+1} \ge \gamma p_k ; k = 1,2,...$$

From lemma 3 it follows that for every sequence  $\{p_k\}$  satisfying H and with  $0 \le p_0 \le \frac{1}{2}$ , there always exists positive 3 and  $\gamma$ . Also by definition of 3, it is clear that for every such sequence  $0 \le 3 \le p_1/p_2$ . Here  $p_2 > 0$ ; in fact because of (12)  $p_k > 0$  for all  $k \ge 1$ . Also  $\gamma$  has to be strictly between 0 and 1. That it cannot be equal to one follows from (18) and the fact that  $\sum_{i=0}^{\infty} p_i$  converges.

The following lemma is the essential lead to the main theorem of this section.

Lemma 4. For every sequence  $\{p_k\}$  with  $0 < p_0 < \frac{1}{2}$  and satisfying H, 3  $\gamma = 1$ .

Proof. From (17) and (18) it is clear that 3  $\gamma \le 1$ . It is sufficient then to prove that 3  $\gamma \ge 1$ . From (14) we have for  $k \ge 2$ ,

(19) 
$$p_k (1-2p_0) = 2 \sum_{k=1}^{\infty} p_i p_{i+k}$$
.

Using (18) on the right side of (19) for each  $p_i$  we have

(20) 
$$p_{k}(1-2p_{0}) \leq \frac{2}{\gamma} \sum_{i=1}^{\infty} p_{i+1}p_{i+k} = \frac{2}{\gamma} \sum_{i=2}^{\infty} p_{i} p_{i+k-1}$$
$$= \frac{1}{\gamma} \lceil p_{k-1} - 2p_{0}p_{k-1} - 2p_{1}p_{k} \rceil,$$

which after simplification, yields for k = 2,3,...,

$$p_{k-1} \ge [\gamma + \frac{2p_1}{1-2p_0}] p_k$$
,

or equivalently for k = 1, 2, ...,

(21) 
$$p_{k} \ge \left[\gamma + \frac{2p_{1}}{1 - 2p_{0}}\right] p_{k+1}.$$

Comparing (17) and (21) and keeping the definition of 3 in mind, we have

(22) 
$$(3 - \gamma) \ge \frac{2p_1}{1 - 2p_0} .$$

Again, using (17) on the right side of (19), we have for  $k \ge 2$ ,

(23) 
$$p_{k}(1-2p_{0}) \geq 2 \beta \sum_{i=1}^{\infty} p_{i+1} p_{i+k} = 8(2 \sum_{i=2}^{\infty} p_{i} p_{i+k-1})$$
$$= \beta \left[p_{k-1} - 2p_{0}p_{k-1} - 2p_{1}p_{k}\right].$$

On simplification, (23) yields for  $k \ge 2$ ,

$$p_{k} \ge \frac{3(1-2p_{0})}{(1-2p_{0}+2p_{1}^{3})} p_{k-1}$$
,

or equivalently for  $k \geq 1$ ,

(24) 
$$p_{k+1} \ge \frac{3(1-2p_0)}{(1-2p_0+2p_1\beta)} p_k.$$

Finally comparing (18) and (24) and using the definition of  $\gamma$ , we obtain

$$\gamma \ge \frac{9(1-2p_0)}{(1-2p_0+2p_1^3)}$$

or after simplification

(25) 
$$(3-\gamma) \le \frac{2p_1}{1-2p_0} 3 \gamma .$$

Now it easily follows from (22) and (25) that 3  $\gamma \ge 1$ .

Q.E.D.

We are now in a position to state and prove the main theorem of this section. 

Theorem 2. Let  $X_1$  and  $X_2$  we two independent copies of a nonnegative discrete random variable X satisfying condition C. Then X and the absolute difference  $|X_1-X_2|$  have the same distribution if and only if the distribution of X is given for some positive constant  $\tau$ , by

(26) 
$$\begin{cases} Pr(X = 0) = p_0 \\ Pr(X = k\tau) = 2p_0(1-p_0) (1-2p_0)^{k-1}; k = 1,2,..., \end{cases}$$

where either  $p_0 = 1 \text{ or } 0 \le p_0 \le \frac{1}{3}$ .

<u>Proof.</u> The case with  $p_0 = 1$  is that of a degenerate r.v. X. Also we have argued before that for a nondegenerate X, we must have  $0 < p_0 \le \frac{1}{2}$ . The case with  $p_0 = \frac{1}{2}$  is covered in theorem 1. Let us assume then that  $0 < p_0 < \frac{1}{2}$ . From lemma 4 and equations (17) and (18) it follows that

(27) 
$$p_{k+1} = \gamma p_k$$
;  $k = 1, 2, ...$ 

or equivalently

(28) 
$$p_{k} = \gamma^{k-1} p_{1} ; k = 1,2,...$$

Now it is easy to show using (9) and the fact that  $\sum_{i=0}^{\infty} p_i = 1$ , that  $\gamma = (1-2p_0)$ 

and 
$$p_1 = 2p_0(1-p_0)$$
. Q.E.D.

ABSOLUTELY CONTINUOUS CASE. Let f(x) denote the p.d.f of the nonnegative r.v. X with the property H. The property H is then equivalent to f(x) satisfying the relations

(29) 
$$\int_{0}^{\infty} f(x)dx = 1 ; f(t) = 2 \int_{0}^{\infty} f(x+t) f(x)dx ; \text{ for all } t \ge 0.$$

Furthermore in view of theorem 1, X is unbounded. The following lemma gives certain properties of an f satisfying (29), which we shall need later.

Lemma 5. Let the p.d.f f(x) of a nonnegative r.v X satisfy (29). Then it also satisfies the following:

- (i) f(x) is lower semicontinuous for all  $x \ge 0$ .
- (ii) f(x) > 0, for all  $x \ge 0$ .

Proof. (i) From (29) for t = 0, we have

(30) 
$$f(0) = 2 \int_0^\infty f^2(x) dx \; ; \int_0^\infty f(x) dx = 1 \; ,$$

so that we must have f(0) > 0. On the other hand since

$$\psi(z) = \int_0^\infty f(y+z) f(y)dy ; -\infty < z < \infty ,$$

is the p.d.f of  $X_1-X_2$ , or equivalently the p.d.f. of the convolution of  $X_1$  and  $-X_2$ ,  $\psi(z)$  is lower semicontinuous for all  $-\infty < z < \infty$ . By virtue of (29) therefore, f(t) is lower semicontinuous for all  $t \ge 0$ .

(ii) Assume that there exists an interval (a,b) with  $\epsilon \le a \le b$ , such that f(x) = 0 for all  $x \in (a,b)$ . Since f(0) > 0 and f(x) is lower semicontinuous at zero, there is an  $\epsilon > 0$ , with  $a \le b - \epsilon/2$ , such that f(x) > 0, for all  $x \in [0,\epsilon]$ . Using this and (29), it is now easy to show that f(z) = 0, for all  $a \le z \le b + \epsilon/2$ , a.e. $\mu$ . By an induction argument we then have f(z) = 0 for all  $a \le z \le b + \epsilon/2$ , a.e. $\mu$ , for  $n = 1,2,\ldots$ . Letting  $n \to \infty$ , we have f(z) = 0 for all  $a \le z \le \infty$ , a.e. $\mu$ . But this implies that X is bounded, which is a contradiction. Thus there exists no interval  $1 \subset [0,\infty)$  with  $\mu(1) > 0$  such that f(x) = 0 for all  $x \in I$ . This implies that f(x) > 0 for all  $x \ge 0$ , a.e. $\mu$ . Now let f(x) = 0 for some f(x) > 0. Then

$$\int_0^{\infty} f(x+x_0) f(x)dx = 0 \Rightarrow \int_a^b f(x+x_0) f(x)dx = 0, \text{ for } 0 < a < b < \infty$$

$$\Rightarrow f(x+x_0) = 0, \text{ for all } a < x < b \text{ with } f(x) > 0, \text{ a.e.}\mu.$$

But  $\mu$  [x: a < x < b, f(x) > 0] = b-a, which also yields  $\mu$  [x+x<sub>0</sub>: a < x < b, f(x) > 0] = b-a > 0. This contradicts the fact that f(x) > 0, for all  $x \ge 0$  a.e. $\mu$ . Thus f(x) > 0, for all  $x \ge 0$ .

We shall now consider a more general problem. Let X and Y be two nonnegative independently but not necessarily identically distributed random variables. Let F and G denote the D.F's of X and Y respectively. Given F and that the distributions of Y and |Y-X| are identical, what can we say about the distribution of Y, i.e. about G? The reader may find in Feller [1] a treatment of this problem considered for a somewhat restricted case. That the distributions of Y and |Y-X| are identical is equivalent to the relation

(31) 
$$G(t) = \int_0^\infty G(x+t) dF(x) + \int_0^\infty F(y+t) dG(y) ; \text{ for all } t \ge 0 .$$

The following theorem provides an answer to the question raised above.

Theorem 3. Let X and Y be two nonnegative independent random variables with F and G as their respective D.F.'s. Let E X <  $\infty$  and F have an absolutely continuous part. Then G satisfies (31) if and only if G is solutely continuous with p.d.f. g where

(32) 
$$g(y) = [1-F(y)] / EX$$
; for all  $y > 0$ .

<u>Proof.</u> It is easy to verify that g(y) of (32) does satisfy (31). All we need to show is that this is the unique g that satisfies (31). To this end, consider a sequence of i.i.d. random variables  $X_1, X_2, X_3, \ldots$ , with their common distribution same as that of X. Define another sequence of random variables  $Z_k$  recursively by

(33) 
$$Z_1 = X_1, Z_{n+1} = |Z_n - X_{n+1}|$$
.

Then clearly  $\{Z_n\}$  is a Markov chain (M.C) with state space  $[0, \infty)$  and the transition D.F. H, given by

$$dH(x|y) = dF(x-y) + dF(x+y),$$

so that if  $G_n$  is the D.F. of  $Z_n$ , it is easily observed that  $G_1 = F$  and for  $n = 2, 3, \ldots$ ,

(35) 
$$G_{n}(y) = \int_{0}^{\infty} G_{n-1}(x+y) dF(x) + \int_{0}^{\infty} F(x+y) dG_{n-1}(x).$$

Letting  $n \to \infty$ , we observe that any solution G of (31) is a stationary distribution of the M.C.  $\{Z_n\}$ . We have already observed that g(y) given by (32) is such as stationary distribution. That this is the unique stationary distribution and hence the unique solution of (31) follows from the fact that the above M.C. defined on  $\{0,\infty\}$  is indecomposable, which is a simple consequence of the fact that F has an absolutely continuous part.

In answer to our original question, we now have the following theorem. Theorem 4. Let  $X_1$  and  $X_2$  be two independent copies of a nonnegative r.v. X with p.d.f. f(x). Then X and  $|X_1-X_2|$  have the same distribution, if and only if for some 0 > 0,

(36) 
$$f(x) = \begin{cases} 0 e^{-\theta x}, & \text{for } x \ge 0 \\ 0 & \text{elsewhere } . \end{cases}$$

<u>Proof.</u> Clearly if f(x) satisfies (36), the distributions of X and  $|X_1-X_2|$  are identical. Assuming now that the distributions of X and  $|X_1-X_2|$  are the same, it is easily seen that f(x) satisfies the conditions of theorem 3, in view of lemmas 1 and 5. On the other hand comparing (29) and (31), we have under H, g(y) = f(y), so that replacing g(y) with f(y) in (32), and solving the resulting equation for  $f(y) = F^1(y)$  we obtain (36) with  $\theta = EX$ . Q.E.D.

5. A FEW CONCLUDING REMARKS. The lines of proof adopted for theorem 3 and hence of theorem 4, in principle should also work for the discrete case of section 3. Let X and Y be two appropriate nonnegative discrete r.v, both independently but not necessarily identically distributed with  $\{p_x\}$  and  $\{p_y\}$  as the set of their probabilities (as defined in section 1). Given that the distributions of Y and  $\|Y-X\|$  are identical, the analogue of equation (29) is given by

(37) 
$$\begin{cases} q_0 = \sum_{x \ge 0} p_x q_x \\ q_y = \sum_{x \ge 0} p_x q_{x+y} + \sum_{x \ge 0} q_x p_{x+y}; \text{ for } y > 0. \end{cases}$$

However, here essentially it is a matter of first guessing a general solution of (37) for  $q_y$ 's satisfying  $\sum q_y = 1$ , in terms of  $p_x$ 's, an analogue of (32). After this, replacing  $q_y$ 's with  $p_x$ 's in this solution,  $p_y$ 's can be explicitly obtained to yield the answer to our original problem.

Concerning the singular case of an X with property H, at present we can only say in view of theorem 1, that X has to be unbounded. On the other hand let us consider again the approach adopted in section 4. Let F(x) and G(y) respectively be the continuous distribution functions of two nonnegative independent random variables X and Y. This will cover both absolutely continuous and singular cases of our problem. Introduce a M.C similar to the one of section 4, defined on  $[0, \circ)$ , but with the assumption that X has the continuous D.F. F(x), so that (31) and (35) are still satisfied. Any solution of (31) is a stationary D.F. G of our M.C.  $\{Z_n\}$ . On the other hand, if  $EX < \infty$ , it is easily verifiable that

(38) 
$$dG(y)/dy = [1-F(x)] / EX; j \ge 0$$
,

is a solution of (31) and hence a stationary D.F. of M.C.  $\{\mathcal{Z}_n\}$ . The only problem

here is to show that (38) is the unique solution of (31). For this we need to show that the M.C.  $\{Z_n\}$  is indecomposable. Once this is established, (38) is the unique solution of (31). The solution to our problem is then obtained by replacing G with F in (38) and solving this for F. This turns out to be the same as (36). Thus, subject to the uniqueness of the solution of (31), the solution to our original problem would be (36) even when D.F. of X is given to be only continuous. This would mean that there is no singular distribution with the property H. Our conjecture is that this is in reality the case.

Finally, in section 3 for the discrete case the result of theorem 2 was proved subject to the condition C. Our conjecture is that this result holds even without this extra condition.

### REFERENCES

<sup>[1]</sup> Feller, W. (1966). An introduction to probability theory and its applications, Vol II, J. Wiley, New York.

<sup>[2]</sup> Puri, Prom S. (1966). Probability generating functions of absolute difference of two random variables, Proc. Nat. Acad. Sciences, 56, No. 4.

Security Classification

DOCUMENT CO	NTROL DATA - R&I		the overall report is classified)					
CORIGINATING ACTIVITY (Corporate author)	•		AT SECURITY CLASSIFICATION					
Purdue University		Unclas	sified					
		26 GROUP						
A Characterization Based on the Absolu Variables.	te Difference o	of Two I	.I.D. Random					
4 DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report, January, 1970		<del>,</del>						
5 AUTHOR(S) (Last name, first name, initial) Puri, Prem S. and Rubin, Herman								
6 REPORT DATE January, 1970	74. YOTAL NO. OF P.	AGES	75 NO. OF REFE					
8. CONTRACT OR BRANT NO. NOOO14-67-A-0226-0008 and GM-10525 b. PROJECT NO.	Se. ORIGINATOR'S REPORT NUMBER(S) Mimeo Series # 218							
c.	3b. OTHER REPORT NO(3) (Any other numbers that may be essigned this report)							
d. 10 AVAILABILITY/LIMITATION NOTICES								
Distribution of this document is unlimited.								
II SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY							
Also supported by the National Institutes of Health	Office of Naval Research Washington, D.C.							
13 ABSTRACT Let X be a nonnegative rando	m variable with	X, and	X <sub>2</sub> as its two indepen-					
dent copies. The problem considered here is to characterize all the nonnegative dis-								
tributions with the property that the distribution of the absolute difference   X -X								
is the same as that of X. It is shown that in general such a distribution has to be								
either purely discrete, or absolute continuous or singular and that it cannot be								
their mixture. It is shown that among lattice distributions the only distribution								
that enjoys the above property is given for some positive constant a, by $P(X = 0) =$								
$p_0$ ; $P(X = k a) = 2p_0(1-p_0) (1-2p_0)^{k-1}$ ; $k = 1,2,3,$ , where either $p_0 = 1$ or								
$p_0 \le \frac{1}{2}$ . When X is absolutely continuous, it is shown that in order to enjoy the								
bove property, its distribution has to be exponential with p.d.f. $f(x) = \theta e^{-\theta x}$ , for $x \ge 0$ , and zero electrons with $\theta \ge 0$ . The case show $x$ is also as								
for $x \ge 0$ , and zero elsewhere, with $\theta > 0$ . The case when X is singular is not completely solved. However, it is shown that in this case X cannot be bounded. A par-								
tial solution has also been given for the following more general question. Let X								
and Y be two nonnegative independently but not necessarily identically distributed and om variables with the property that the distributions of Y and $ Y-X $ are identical. Given the distribution of X, what an we say about the distribution of Y?								
			أحسب والمراجع					

### Unclassified

Security Classification

14 KEY WORDS		LINK A		LINK II		LINKC	
	KEY WORDS	ROLC	WT	HQU.L	0.7	HOLE	W T
1.	Characterization of distributions.						1
2.	Absolute difference of two nonnegative random variables.						
3.	Exponential distribution.						
	·	-					

#### INSTRUCTIONS

- ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Delense activity or other organization (corporate author) issuing the report.
- 2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. GROUP: Automatic downgrading is specified in DoD Directive \$200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningfut title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is convered.
- 5. AUTHOR(S): Enter the name(s) of author(s) on shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
- e. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7.: DOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- $76-{\rm NUMBER\,OF\,REFERENCES}$  . Enter the total number of references cuted in the report.
- No. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 86. W. & 8d. PROJECT NUMBER: Enter the appropriate multitary department identification, such as project number, subjectly to number, system numbers, task number, etc.
- 9n ORIGINATOR'S REPORT NUMBER(S): Enter the offitial report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- on OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).
- to. AVAILABILITY, LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

- 11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include adgress.
- 13. ABSTRACT: Enter an abatract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U)

There is no limitation on the length of the obstruct. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cutaloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.